

Ripple and kink dynamics

H. Caps and N. Vandewalle
*GRASP, Institut de Physique B5, Université de Liège,
 B-4000 Liège, Belgium.*

We propose a relevant modification of the Nishimori-Ouchi model [*Phys. Rev. Lett.* **71**, 197 (1993)] for granular landscape erosion. We explicitly introduce a new parameter: the angle of repose θ_r , and a new process: avalanches. We show that the θ_r parameter leads to an asymmetry of the ripples, as observed in natural patterns. The temporal evolution of the maximum ripple height h_{max} is limited and not linear, according to recent observations. The ripple symmetry and the kink dynamics are studied and discussed.

(Accepted for publication in *Phys. Rev. E*)

45.70.Mg, 81.05.Rm, 64.75.+g

I. INTRODUCTION

The formation of ripples due to the wind blowing across a sand bed [1] is a complex phenomenon involving various mechanisms such as air turbulence, non-linear fluxes of sand, saltation, grain ejection threshold, creeping and avalanches.

Different models [2–5] for ripple formation have been proposed. Some of those models consider both hopping and rolling of grains, and they are able to reproduce travelling ripple structures. Simulations [6] have also considered various additional effects like the screening of crests, the grain reptation and the existence of a grain ejection threshold.

The Nishimori-Ouchi (NO) model [3] is able to reproduce a wide variety of different eolian structures: ripples, star dunes, barchanic dunes, etc... The main advantage of such a numerical model is that all parameters can be easily tuned and the dynamics can be deeply investigated. In a recent work [7], we have shown that the NO model reproduces the complex labyrinthic patterns created in some experiments [8]. However, the NO model leads to unrealistic features. Among others, a cube of sand can be stable within the NO model. It is possible to create aeolian structures with an infinite height. In fact, the existence of a critical angle is lacking in the NO model.

In the next section, we will propose a modification of the NO model in order to impose a more realistic constraint to granular surfaces. In section 3, we will study how the ripple formation depends on the various parameters of the model. In section 4, the dynamics of the new model will be studied and will be confronted to various situations. Finally, a summary of our findings will be given in section 5.

II. MODELS

In this section, we present two 2-dimensional models of granular landscape evolution.

A. Nishimori-Ouchi (NO)

In the Nishimori-Ouchi model [3], two modes of granular transport are considered: (i) saltation and (ii) reptation. The saltation process acts along the x axis. It takes grains from a site $(x - \ell, y)$ and add these grains to a site (x, y) . The value of the *saltation length* $\ell(x, y)$ depends on the local slope of the landscape at (x, y) . In this model, the saltation grains are captured by the surface when they touch it. The granular flux Q_{salt} due to the saltation process can be written as

$$Q_{salt} = A \int_{x-\ell}^x N(x', y) dx' , \quad (1)$$

where $N(x, y)$ is the amount of sand per time unit and surface unit at the position (x, y) . A is a dimensionless constant parameter.

The NO model also considers a 2-dimensional diffusion process, i.e. the *reptation*. The amount of sand displaced at (x, y) is supposed to be proportional to the local curve of the surface at that position. The flux Q_{rept} due to the reptation reads

$$Q_{rept} = D \nabla h, \quad (2)$$

with a kind of diffusion coefficient D . The mass conservation imposes

$$\frac{\partial h}{\partial t} = -\nabla Q, \quad (3)$$

where $Q = Q_{salt} + Q_{rept}$. Therefore, the temporal evolution of the height h of sand at the position (x, y) reads

$$\begin{aligned} \frac{\partial h(x, y, t)}{\partial t} = & A \left[N(x - \ell, y) \left(1 - \frac{\partial \ell(x, y)}{\partial x} \right) - N(x, y) \right] \\ & + D \Delta h. \end{aligned} \quad (4)$$

As in the NO model [3], we have assumed that the local slope (along the x direction) mainly controls the sand flux Q_{salt} . Experimentally [1], one observes that the wind is stronger on the crests. As a consequence, the amount of displaced sand is larger around the crests. This effect can be included in the NO model by assuming

$$\frac{\partial Q_{salt}}{\partial x} = \delta \left(1 + \tanh \frac{\partial h}{\partial x} \right) \left(1 + \varepsilon - \tanh \frac{\partial h}{\partial x} \right) \quad (5)$$

where the parameters δ and ε give respectively the scale factor of the saltation flux and the non-zero asymptotic quantity of sand displaced when $\partial h / \partial x > 0$. The saltation length ℓ is represented by the factor $(\tanh(\frac{\partial h}{\partial x}) + 1)$. This assumes that ℓ is maximum on the crest and that small saltation jumps take place on the downstream part of the ripple.

We have numerically solved the NO model on two-dimensionnal square lattices starting from an initially plane landscape. At each time step, a randomly chosen site is submitted to both saltation and reptation processes (Eqs.(4) and (5)). Figure 1 shows the maximum height h_{max} reached by the profile as a function of time t . Time is expressed in numerical steps (iterations).




FIG. 1. Evolution of the maximum height h_{max} of the surface as a function of time t in the case of the NO model. The simulation parameters are : $\delta = 25$, $D = 0.4$ and $\varepsilon = 0.3$. The lattice size is 101×101 .

One observes that : (i) the growth of the maximum height h_{max} is linear and (ii) there is no saturation. This means that ripples become infinitely high! This unrealistic behavior comes from the right hand side of Eq.(4). Indeed, when the first term becomes larger, the sand deposit becomes more important than the relaxation. As time goes on, a net growth of the profile is thus recorded. This situation will remain unchanged since the ratio between deposit and relaxation is fixed once by the values of δ , ε and D . Consequently, h_{max} grows *ad infinitum*. One should note that this pathology implies the possibility of making cubes of sand since nothing prohibits vertical walls. In order to get a more realistic situation, an additional term should be included in the NO model.

B. Saltation-Creep-Avalanche (SCA)

Granular materials are characterized by the *angle of repose* θ_r [9]. If the slope of a granular pile exceeds $\tan(\theta_r)$, an avalanche is created and stabilizes the surface. The angle of repose depends on the grain characteristics: size distribution, shape, rugosity, and density. Typically, this angle θ_r is about 37° for sand [1,9]. Since it is an intrinsic property of the granular material, the relevant parameter θ_r should be considered in numerical models.

Let us consider a square lattice with discrete spatial variables x and y . The stability condition with respect to the angle of repose θ_r reads

$$\begin{aligned} |h_x - h_{x+1}| &< \tan(\theta_r) \\ |h_y - h_{y+1}| &< \tan(\theta_r) \end{aligned} \quad (6)$$

where the left hand side is the discrete counterpart of the spatial derivative of the height of sand h at the position (x, y) . If Eq.(6) is not verified for a specific direction (x or y), an avalanche occurs as illustrated in Figure 2. In this case, the excess of sand $E = \tan(\theta) - \tan(\theta_r)$ (with $\tan(\theta) = \frac{\partial h}{\partial i}$, with $i = x$ or $i = y$ depending on which condition (6) is not verified) is supposed to be equitably distributed between x and $x + 1$ if the avalanche occurs in that direction. The same applies for the y direction. Taking those considerations into account, Eq.(4) becomes

$$\begin{aligned} \frac{\partial h(x, y, t)}{\partial t} = A \left[N(x - \ell, y) \left(1 - \frac{\partial \ell(x, y)}{\partial x} \right) - N(x, y) \right] \\ + \Theta(E) \frac{E}{2} + D \Delta h, \end{aligned} \quad (7)$$

where Θ is the Heavyside step function and

$$E = |\nabla h| - \tan(\theta_r). \quad (8)$$

The new model is called the Saltation-Creep-Avalanche (SCA) model. It will be investigated in the present paper and numerical results will be compared with respect to NO landscapes. One should note that we have numerically solved Eq.(7) using the same flux law (5) as in the original NO model.




FIG. 2. Avalanches lead to a profile stabilisation: (left) the local slope $\tan(\theta)$ is larger than the critical value $\tan(\theta_r)$, the surface is unstable. (right) The avalanche brings the local slope to a new stable value $\tan(\theta_r)$. Note that $E = |\nabla h| - \tan(\theta_r)$.

III. RIPPLE FORMATION

In this section, we investigate the influence of the four parameters δ , ε , D and θ_r on the formation of the ripples in the SCA model. All the simulations presented herebelow are made on 150×150 square lattices. The evolution of the landscape is typically stopped after 10^7 iterations. The default parameter values are $\delta = 12.5$, $\varepsilon = 0.3$, $D = 0.4$ and $\theta_r = 37^\circ$.

A. Flux parameter δ

The saltation process obeys Eq.(5) in which the parameter δ controls the flux Q_{salt} . A small arbitrary value of δ implies that small amounts of sand are displaced along small saltation lengths, while larger values of δ lead to a more important transport of sand over long distances. Experimentally [1], one observes that the ripple wavelength (the distance between two consecutive crests) grows with the saltation length ℓ .

We have performed simulations varying δ from 1 to 100. For very small values of the parameter ($\delta < 7$), no ripple appears and the surface remains irregular. This threshold is not encountered in the NO model and is certainly due to the angle of repose. For small values of the flux parameter, not enough sand is displaced and the surface is continuously smoothed by avalanches.

Above this threshold $\delta \simeq 7$, a set of several ripples appears. The density of ripples depends on δ . The larger δ is, the smaller is the density of ripples. For example, after $5 \cdot 10^6$ iterations, the density of ripples is 0.04 for $\delta = 10$, while there is only a single ripple (density of 0.006) if $\delta = 85$. The SCA model is thus in good agreement with observations: the wavelength of the ripple pattern grows with the saltation length ℓ . In the case of the NO model, the density of ripples grows non-linearly with δ in opposition with experimental observations.

B. Asymmetry parameter ε

In the SCA model, we have assumed a non-zero value for the flux of sand Q_{salt} displaced on the stoss slopes (which are upstream) of the ripples (see Eq.(5)). This hypothesis is supported by numerical results [3]. If the parameter ε is non zero, the saltation flux becomes asymmetric (see Figure 3). Sand is always taken from the stoss slope, while Q_{salt} tends to zero on the lee slope. This comes from the fact that sand on the lee slope is confined in wind vortices.




FIG. 3. The normalized saltation flux Q_{salt}/δ of Eq.(5) as a function of the spatial derivative ∇h of the height of sand h . Different values of the parameter ε are illustrated.

Practically, we have considered values of ε ranging from 0 to 2. Our observations are: (i) no ripple appears if $\varepsilon = 0$. (ii) for small values of ε (typically $0 < \varepsilon < 0.2$), undulations appear in the landscape. (iii) As ε becomes larger, the number of ripples present in the system grows (from 5 at $\varepsilon = 0.5$ to 11 for $\varepsilon = 1.8$). Moreover, ripples are more asymmetric (windward slopes smaller than downstream slopes) for large values of ε . We will study this asymmetry in section 4.

In the case of the NO model we observe that: (i) if $\varepsilon = 0$ very small ripples appear. (ii) when the parameter reaches a threshold $\varepsilon \simeq 0.2$ larger ripples appear. Note that the wavelength of the ripples does not depend on ε above that threshold. The fact that ripples appear for $\varepsilon = 0$ in the NO model and do not appear within the SCA model is due to the existence of an angle of repose. Indeed, avalanches tend to smooth the landscape.

In summary, the flux Q_{salt} should be asymmetric in order to create ripples. For positive slopes, a minimum value of this flux is required.

C. Reptation coefficient D

The parameter D plays the role of a diffusion coefficient for smoothing the irregularities along the surface. We have considered values of D ranging from 0 to 1. Once again, no ripple appears for very small values of that parameter. Ripple appearance is allowed in the interval $0.1 \leq D \leq 0.5$. If $D < 0.1$ the surface remains irregular. There is nearly no interaction between the cells because of the small diffusion coefficient. Note that the same behavior is observed in the NO model. As D becomes larger, the ripple wavelength increases. Finally, for $D > 0.5$ the diffusion coefficient becomes too large and the surface remains smooth and flat. For the NO model, no ripple appears when $D > 1.8$, a larger value than in the case of the SCA model.

D. Angle of repose θ_r

Parameters D and θ_r are intrinsic parameters of the granular material, while δ and ε should be related to the wind. At each iteration, the reptation acts and moves grains. On the other hand, an avalanche occurs if and only if the local slope of the surface exceeds the angle of repose. The role played by both D and θ_r parameters is thus different: D is a diffusion coefficient, which controls the flow of sand between neighboring cells, while the angle of repose θ_r governs the local slope of the granular surface.

We considered typical values of θ_r between 20° and 40° . For ripples of the same wavelength, we have noted that small angles of repose imply small ripple heights. The angle of repose is thus responsible of the value of the ratio h_{max}/λ , where λ is the ripple wavelength. Indeed, a slope larger than $\tan(\theta_r)$ is reduced to a value $\tan(\theta_r)$ by avalanches. We have also noted the appearance of ripples for all tested values of θ_r .

IV. RIPPLE DYNAMICS

In this section, we investigate the dynamics of the SCA model. Three different aspects will be considered: the ripple height, the ripple shape and the formation of kinks. Again, the default parameters values are $\delta = 12.5$, $\varepsilon = 0.3$, $D = 0.4$ and $\theta_r = 37^\circ$.

A. Ripple amplitude h_{max}

Figure 4 presents the temporal evolution of the maximum ripple height h_{max} in the case of the SCA model. One should note the saturation of h_{max} for long times. This figure should be compared with Figure 1.




FIG. 4. Evolution of the maximum height h_{max} of the landscape as a function of time t in the case of the SCA model. The data have been fitted using Eq.(9). The fit parameters are $a = 230.8 \pm 0.6$, $b = 377.3 \pm 3$ and $c = 1.33 \cdot 10^7 \pm 0.01 \cdot 10^7$. The simulation parameters are the same as those of Figure 1. The angle of repose is $\theta_r = 37^\circ$. When all ripples have merged, the system contains only one ripple for which the height is limited by the angle of repose.

In Figure 4, many gaps can be observed in the early stages of evolution. Those gaps come from the absorption of small ripples by the largest ones. Since a small ripple travels faster than a large one [1], the large ripple tends to absorb the small. Figure 5 presents the evolution of a part of the lattice for different stages of the simulation. For each picture, a transverse view of the landscape is shown. One should note that the merge occurs as follows: first, the small ripple climbs on the larger one. The arrival of the small ripple on the crest causes an avalanche. A part of the small ripple falls in the avalanche, while the remainder feeds the larger one. It results in a fast growth of the largest ripple even it moves slowly.




FIG. 5. The merge of two ripples. The wind blows from left to right. The profile of the landscape is shown below each picture. (*top row*) the smallest ripple travels faster than the largest one. (*second row*) The small ripple climbs on the large one. (*third row*) An avalanche is initiated by the arrival of the small ripple on the crest. (*bottom row*) The small ripple ‘feeds’ the large one. Note that the profiles have been rescaled in order to be compared.

Asymptotically, only one ripple occupies the whole lattice, and the time needed for a global merge becomes larger as δ decreases. If δ is large, the initial density of ripples is small. In this case, the global merge is obtained after a small number of collapses. In opposition, a large number of merges is necessary for small values of δ . If one assumes that ripple collapses is a size independent process, one can understand that the global merge is faster when $\delta \gg 1$.

The main characteristics of $h_{max}(t)$ are (i) gaps in the early stages of ripple formation, and (ii) a saturation for long times. Looking for details in Figure 4, one could see that gaps are allways followed by the same kind of growth. Indeed, the maximum ripple height evolves according to an exponential growth law

$$h_{max} = a - b \exp\left(\frac{-t}{c}\right), \quad (9)$$

where a , b and c are fitting parameters. This law is shown in Figure 4. One should see that the fit has been performed after the primary gaps, e.g. for $t > 10^7$ iterations.

In summary, we have seen that, contrary to the NO model, the SCA model naturally leads to a non-linear evolution of h_{max} . Note that this kind of behavior has been experimentally observed in [10,11]. Moreover, the SCA model predicts a saturation of h_{max} after a finite time, as expected and observed. The saturation value depends essentially on θ_r . The larger is θ_r , the higher are the ripples.

B. Ripple aspect ratio σ

Under a non-oscillating wind, a ripple has generally an asymmetric shape [1,8]. The stoss slope is indeed smaller than the lee slope. In Figure 6, the profile of the ripples is shown for both NO and SCA models. Ripples are more symmetric in the NO model than in the SCA model.




FIG. 6. Height h of sand (in arbitrary units) as a function of the spatial coordinate x . Both NO and SCA models are shown. The curves have been rescaled in order to be compared.

In order to measure the aspect ratio σ of a ripple, we have measured the ratio of the horizontal projections of both sides of the ripples. Let us call x_s the projection of the stoss slope on the horizontal x axis, and x_l the projection of the lee slope. The aspect ratio of a ripple is defined as

$$\sigma = \frac{x_s}{x_l}. \quad (10)$$

The measure of σ is averaged over all the ripples of the landscape. If $\sigma = 1$ the ripples are symmetric, while $\sigma \neq 1$ implies an asymmetry.

In Figure 7, the temporal evolution of σ for both NO and SCA models are reported. Typical values of σ range between 1 and 5 in the SCA model. In the NO model, σ remains close to 1.

In the case of the SCA model, one should note that σ grows linearly with time, and then remains close to a saturation value. The appearance of a saturation seems coherent since only one ripple occupies the whole lattice after a finite time. The angle of repose limits the height of that ripple and its morphology.




FIG. 7. Temporal evolution of the aspect ratio σ . Both NO and SCA models are illustrated. The parameter values are $\delta = 12$, $\varepsilon = 0.3$, $D = 0.4$ and $\theta_r = 37^\circ$.

In order to compare the four parameter dependencies on σ , we have considered a reference situation which corresponds to the default parameters. Varying the value of a parameter while keeping constant the others allows us to define the influence of that parameter on the temporal evolution of the aspect ratio. Herebelow, we describe the results obtained for each of the four parameters δ , ε , D and θ_r . One should note that in all cases, the temporal evolution of σ begins with a quasi-linear law, followed by a saturation.

The aspect ratio increases with the parameter δ . Since the ripple wavelength directly depends on δ , that result means that small ripples are less asymmetric than large ones. This result is in good agreement with observations [1,8].

Figure 8 presents the temporal evolution of the aspect ratio σ for different values of ε . One can see that small

ripples (for small ε values) are less asymmetric. Indeed, σ is larger for $\varepsilon = 1$ than for $\varepsilon = 0.3$. We have seen in Section III.B. that the ripple wavelength becomes larger as ε increases. One should also note that σ saturates more rapidly when ε is small. This is consistent with the fact that the time needed for a global merge is shorter in the case of large ripples.

As observed for δ and ε , the aspect ratio follows any variation of D . This results is in agreement with those observed previously in this section. Indeed, we have seen in Section III.D. that the ripple wavelength increases as D becomes larger.

We have taken realistic values of θ_r between 30° and 40° . The aspect ratio σ is found to be independent of the value of θ_r . This result is outside the scope of this paper and will be studied in the future.




FIG. 8. Temporal evolution of the aspect ratio σ in the case of the SCA model. Different values of ε are illustrated.

C. Kink dynamics

Although the wind blows in a single direction, the resulting ripples are not strictly perpendicular to the wind direction. There are defects. Two different classes of defects can be found: *kinks* which are fusion points of two crests, and *antikinks* which are endings of crests. In Ref. [7], we have studied in the NO model how defects modify a landscape morphology when a brutal change occurs in the wind direction. We have also shown in Ref. [7] that under a constant wind direction, the density of defects decreases as time goes on. In the present section, we will focus on the processes leading to this decrease in the SCA model.

After extensive simulations, we conclude that some kinks can propagate during very long times, while others disappear quite rapidly.

The disappearance of a defect mainly occurs when a ripple is splitted into two inequitably parts (see an example in Figure 9). The small branch moves faster than the larger one and climbs on it following the merging dynamics described in section IV.A. Reaching the crest, the

small branch creates an avalanche. The sand rolling on the lee slope may be captured. If the amount of sand escaped from this avalanche is small, the defect disappears and the small branch vanishes. After this process, the number of defects is reduced. If all defects show this kind of behavior the defect density will rapidly tend to zero. However, this is not the case [7].

The propagation of a kink takes place when a sufficient amount of sand escapes from the avalanche. This is the case when the branches are quite long. In the emphasized part of Figure 10, one can see the ending of a small ripple. The extremity of the ripple moves faster than the main part. The merge (second image) is accompanied by an avalanche. As seen in the previous paragraph, sand is expelled out of the avalanche. But here, the amount of sand is sufficient to create a new ripple. The merging/propagation process is thus repeated on the next ripple.

Let us consider a branch of length ℓ_b . The merge of this branch with a larger ripple causes the decrease of the branch length since a part of the branch has been owned by the ripple. When a second meeting occurs, ℓ_b is once again reduced. As a consequence, the initial length of the branch is a relevant parameter in order to predict the life-time of a defect. For example, a kink of initial length $\ell_b = 65$ can typically propagate during 9.5×10^5 iterations, while a kink of initial length $\ell_b = 23$ has a life expectancy of 2.5×10^5 iterations. One should note that those life-times mainly depend on the diffusion coefficient D and the avalanche process.

In Ref. [7], we have found a non zero asymptotic value for the defect density. The existence of propagating kinks gives a picture of this behavior. The relevant ingredient for kink disappearance being the length of the branch.




FIG. 9. The disappearance of a kink. (top row) The smallest branch of the ripple climbs on the main part. (bottom row) the avalanche feeds the ripple, and the branch vanishes.




FIG. 10. The propagation of a kink. The defect is expelled downstream.

V. SUMMARY

We have introduced new ingredients in the Nishimori-Ouchi (NO) model: the existence of an angle of repose θ_r and subsequent avalanches when the local slope θ is larger than θ_r . The new model reproduces realistic features of aeolian ripples such as a non-linear evolution of the ripple amplitude h_{max} . The origin of this behavior has been explained by the merge of ripples traveling at different speeds. Increasing the saltation length, we have observed a growth of the ripple wavelength, in agreement with observations. In the case of a constant wind orientation, natural ripples are asymmetric. This feature has also been reproduced by the new model.

Studying the kink dynamics, we have shown the coexistence of two kinds of kink evolution: propagation and disappearance. The defect initial length has been proposed as a relevant parameter in order to predict the life-time of that defect. From this result, the existence of a non-zero asymptotic value of the kink density has been explained.

ACKNOWLEDGEMENTS

HC is financially supported by the FRIA, Belgium. This work is also supported by the Belgian Royal Academy of Sciences through the Ochs-Lefebvre prize.

- [2] R.B. Hoyle and A.W. Woods, *Phys. Rev. E* **56**, 6861 (1997)
- [3] H. Nishimori and N. Ouchi, *Phys. Rev. Lett.* **71**, 197 (1993)
- [4] N. Vandewalle and S. Galam, *Int. J. Mod. Phys. C* **10**, 1071 (1999)
- [5] W. Landry and B.T. Werner, *Physica D* **77**, 238 (1994)
- [6] R.S. Anderson, *Earth-Sci. Rev.* **29**, 77 (1990)
- [7] H. Caps and N. Vandewalle, submitted for publication (2001).
- [8] D. Goossens, *Earth. Surf. Processes and Landforms* **16**, 689 (1991)
- [9] J. Duran, *Sand, powders and grains*, (Springer, 1997)
- [10] A. Betat, V. Frete, and I. Rheberg, *Phys. Rev. Lett.* **83**, 88 (1999)
- [11] A. Stegner and J.E. Wesfreid, *Phys. Rev. E* **60**, R3487 (1999)

[1] R.A. Bagnold, *The physics of blown sand and desert dunes*, (Chapman and Hall, London, 1941)